

Duration of Untreated Psychosis in First-Episode Psychosis is not Associated With Common Genetic Variants for Major Psychiatric Conditions: Results From the Multi-Center EU-GEI Study

Olesya Ajnakina^{*1-3,6}, Victoria Rodriguez⁴, Diego Quattrone⁵, Marta di Forti⁵, Evangelos Vassos^{5,6}, Celso Arango⁶, Domenico Berardi⁷, Miguel Bernardo⁸, Julio Bobes⁹, Lieuwe de Haan¹⁰, Cristina Marta Del-Ben¹¹, Charlotte Gayer-Anderson¹², Hannah E. Jongsma¹³⁻¹⁵, Antonio Lasalvia¹⁶, Sarah Tosato¹⁶, Pierre-Michel Llorca¹⁷, Paulo Rossi Menezes¹⁸, Bart P. Rutten¹⁹, Jose Luis Santos²⁰, Julio Sanjuán²¹, Jean-Paul Selten²², Andrei Szöke²³, Ilaria Tarricone²⁴, Giuseppe D'Andrea⁷, Alexander Richards^{24,6}, Andrea Tortelli²⁵, Eva Velthorst^{26,27}, Peter B. Jones^{28,29}, Manuel Arrojo Romero³⁰, Caterina La Cascia³¹, James B. Kirkbride^{32,6}, Jim van Os^{4,33,34}, Mick O'Donovan³⁵, and Robin M. Murray^{4,36}; EU-GEI WP2 Group[†]

¹Department of Biostatistics & Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, University of London, London, UK; ²Department of Behavioural Science and Health, Institute of Epidemiology and Health Care, University College London, London, UK; ³Department of Clinical Medicine, Aarhus University, Aarhus, Denmark; ⁴Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK; ⁵Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK; ⁶Child and Adolescent Psychiatry Department, Institute of Psychiatry and Mental Health, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, Madrid, Spain; ⁷Department of Biomedical and Neuromotor Sciences, Psychiatry Unit, Alma Mater Studiorum Università di Bologna, Bologna, Italy; ⁸Department of Psychiatry, Barcelona Clinic Schizophrenia Unit, Neuroscience Institute, Hospital Clinic of Barcelona, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Spain; ⁹Faculty of Medicine and Health Sciences – Psychiatry, Universidad de Oviedo, ISPA, INEUROPA, CIBERSAM, Oviedo, Spain; ¹⁰Department of Psychiatry, Early Psychosis Section, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands; ¹¹Neuroscience and Behavior Department, Ribeirão Preto Medical School, University of São Paulo, São Paulo, Brazil; ¹²Department of Health Service and Population Research, Institute of Psychiatry, Psychology and Neuroscience, King's College London, De Crespigny Park, Denmark Hill, London, UK; ¹³Centre for Longitudinal Studies, University College London, London, UK; ¹⁴Centre for Transcultural Psychiatry Veldzicht, Balkbrug, The Netherlands; ¹⁵University Centre for Psychiatry, University Medical Centre Groningen, Groningen, The Netherlands; ¹⁶Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement Sciences, University of Verona, Verona, Italy; ¹⁷Université Clermont Auvergne, CMP-B CHU, CNRS, Clermont Auvergne INP, Institut Pascal, Clermont-Ferrand, France; ¹⁸Department of Preventative Medicine, Faculdade de Medicina FMUSP, University of São Paulo, São Paulo, Brazil; ¹⁹Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, South Limburg Mental Health Research and Teaching Network, Maastricht University Medical Centre, Maastricht, The Netherlands; ²⁰Department of Psychiatry, Servicio de Psiquiatría Hospital “Virgen de la Luz,” Cuenca, Spain; ²¹Department of Psychiatry, Hospital Clínico Universitario de Valencia, INCLIVA, CIBERSAM, School of Medicine, Universidad de Valencia, Valencia, Spain; ²²Rivierduinen Institute for Mental Health Care, Sandifortdreef 19, 2333 ZZ Leiden, The Netherlands; ²³Univ Paris Est Creteil, INSERM, IMRB, AP-HP, Hôpitaux Universitaires “H. Mondor,” DMU IMPACT, Fondation FondaMental, Creteil, France; ²⁴Division of Psychological Medicine and Clinical Neurosciences, Cardiff, UK; ²⁵Etablissement Public de Santé Maison Blanche, Paris, France; ²⁶Department of Psychiatry, Early Psychosis Section, Academic Medical Centre, University of Amsterdam, Amsterdam, The Netherlands; ²⁷Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY; ²⁸Department of Psychiatry, University of Cambridge, Herchel Smith Building for Brain & Mind Sciences, Cambridge, UK; ²⁹CAMEO Early Intervention Service, Cambridgeshire & Peterborough NHS Foundation Trust, Cambridge, UK; ³⁰Department of Psychiatry, Psychiatric Genetic Group, Instituto de Investigación Sanitaria de Santiago de Compostela, Complejo Hospitalario Universitario de Santiago de Compostela, Santiago de Compostela, Spain; ³¹Department of Experimental Biomedicine and Clinical Neuroscience, University of Palermo, Palermo, Italy; ³²PsyLife Group, Division of Psychiatry, University College London, London, UK; ³³Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, South Limburg Mental Health Research and Teaching Network, Maastricht University Medical Centre, Maastricht, The Netherlands; ³⁴UMC Utrecht Brain Centre, Utrecht University Medical Centre, Utrecht, The Netherlands; ³⁵Division of Psychological Medicine and Clinical Neurosciences, MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, Cardiff, UK; ³⁶Department of Psychiatry, Experimental Biomedicine and Clinical Neuroscience, University of Palermo, Palermo, Italy

*To whom correspondence should be addressed; Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, PO20, 16 De Crespigny Park, London SE5 8AF, UK; tel: +44(0)20 7848 0938, e-mail: olesya.ajnakina@kcl.ac.uk

[†]EU-GEI collaborators and their affiliations are listed in the Acknowledgments section.

Duration of untreated psychosis (DUP) is associated with clinical outcomes in people with a diagnosis of first-episode psychosis (FEP), but factors associated with length of DUP are still poorly understood. Aiming to obtain insights into the possible biological impact on DUP, we report genetic analyses of a large multi-center phenotypically well-defined sample encompassing individuals with a diagnosis of FEP recruited from 6 countries spanning 17 research sites, as part of the European Network of National Schizophrenia Networks Studying Gene-Environment Interactions (EU-GEI) study. Genetic propensity was measured using polygenic scores for schizophrenia (SZ-PGS), bipolar disorder (BD-PGS), major depressive disorder (MDD-PGS), and intelligence (IQ-PGS), which were calculated based on the results from the most recent genome-wide association meta-analyses. Following imputation for missing data and log transformation of DUP to handle skewedness, the association between DUP and polygenic scores (PGS), adjusting for important confounders, was investigated with multivariable linear regression models. The sample comprised 619 individuals with a diagnosis of FEP disorders with a median age at first contact of 29.0 years (interquartile range [IQR] = 22.0–38.0). The median length of DUP in the sample was 10.1 weeks (IQR = 3.8–30.8). One SD increases in SZ-PGS, BD-PGS, MDD-PGS or IQ-PGS were not significantly associated with the length of DUP. Our results suggest that genetic variation does not contribute to the DUP in patients with a diagnosis of FEP disorders.

Key words: polygenic scores/schizophrenia/psychosis/genome-wide association studies/duration of untreated psychosis

Introduction

Despite historical pessimism about schizophrenia prognosis,¹ it has now been recognized that interventions at the onset of first-episode psychosis (FEP), which is an umbrella term used to refer to schizophrenia spectrum disorders or related psychotic disorders, can improve subsequent illness outcomes.^{2,3} This recognition has led to development of early intervention services, which are founded on an assumption that duration of untreated psychosis (DUP), defined as the time from manifestation of first psychotic symptoms to initiation of adequate treatment,⁴ influences treatment outcomes.^{3–5} Despite the widespread introduction of early intervention services, however, individuals suffering with FEP still experience delays of approximately 1–2 years between onset of first psychotic symptoms and initiation of treatment,⁶ prompting fears of serious consequences on patients' lives, including enduring deficits and disability.⁶ It is, of course, possible that the relationship between DUP and psychosis outcomes may be a product of other factors^{7,8} related to the organization of mental health system,

treatment-seeking behaviors, quality of available treatment,⁹ or poor premorbid functioning. Seen in this way, DUP may be a marker of the illness severity rather than a predictor of the illness itself.

Schizophrenia is a highly heritable disorder, with twin studies estimating its heritability to be more than 75%.¹⁰ Genomic studies revealed that the genetic architecture of schizophrenia comprises multiple common risk alleles scattered across the whole genome.¹¹ Built on the results from the genomic studies, polygenic scores (PGS) analyses confirmed that schizophrenia is highly polygenic nature,^{10,11} where its onset is influenced by many common genetic variants of small effects.^{12–14} Further evidence highlighted that the impact of the combined effect of common genetic markers for schizophrenia, as measured with polygenic score for schizophrenia, extends beyond schizophrenia diagnosis. Indeed, a higher polygenic score for schizophrenia was shown to associate with more severe negative symptoms¹⁵; whereas, longer DUP is also associated with more severe negative symptoms at first presentation.¹⁶ Considering negative symptoms were linked to cognitive impairments and deficiencies in social and occupational domains in people with a diagnosis of schizophrenia,² all of which contribute to prolonged delay in seeking help,¹⁷ it is feasible that length of DUP might be influenced by genetic factors.⁸ However, this question has not been investigated.

Because the etiology of FEP is highly multifactorial, it is likely that other factors may have an important impact on the delay between onset of first psychotic symptoms and initiation of adequate treatment. Certainly, the length of DUP were shown to be influenced by reduced cognitive functioning or intelligence,¹⁸ severity of depressive symptoms reported in patients with a diagnosis of FEP disorders¹⁹ and bipolar disorders.^{20,21} Similar to schizophrenia, PGS analyses showed that major depressive disorder, bipolar disorders, and cognition are highly polygenic in nature,^{22–24} with an overlapping, though to varying degree, genetic underpinnings. For example, PGS that combined the additive effect of common genetic markers associated with bipolar disorders discriminated individuals with a diagnosis of schizophrenia²⁵ and major depressive disorder from healthy controls.^{10,25} Although much uncertainty remains about their ultimate clinical utility,²⁶ PGS have the power to considerably advance our knowledge of the underlying nature of complex phenotypes.^{27–29}

Therefore, aiming to obtain insights into possible origins of DUP in people with a diagnosis of FEP disorders, we investigated associations between DUP and PGSs for schizophrenia, bipolar disorders, major depressive disorder, and cognition in a large multi-center phenotypically well-defined sample of individuals with a diagnosis of FEP disorders. Because the length of DUP in individuals with a diagnosis of schizophrenia spectrum disorders is reported to be considerably

longer compared to other psychotic disorders,^{17,30} we additionally investigated if our findings were applicable to all individuals with FEP disorders or were specific to patients with first-episode schizophrenia spectrum disorders. We hypothesized that there will be a positive association between polygenic propensity for schizophrenia, bipolar disorder, major depression, and intelligence with longer DUP in participants with a diagnosis of FEP disorders.

Methods

Sample

Participants were recruited and assessed as part of the incidence and first episode case-control study, conducted as part of the European network of national schizophrenia networks investigating Gene-Environment Interactions (EU-GEI) study.³¹ EUGEI study was designed to investigate risk factors for psychotic disorders between May 2010 and April 2015 in tightly defined catchment areas in 17 sites across 6 countries, which were the United Kingdom, The Netherlands, France, Spain, Italy, and Brazil.³² The research sites within each country were purposefully selected to include a mix of urban and rural areas.^{31,32} The inclusion criteria for FEP cases were: (1) presentation with a clinical diagnosis for a FEP as defined by International Statistical Classification of Diseases and Related Health Problems, 10th Revision (ICD-10) criteria³¹ (codes F20-F33) within the timeframe of the study; (2) aged between 18 and 64 years (inclusive); and (3) resident within one of the 17 defined catchment area at the time of their first presentation to psychiatric services for psychosis. Because the construction of PGSs is dependent on the availability of the summary statistics from genome-wide association studies (GWASs), which are currently based on population of European descent,³³ for the purpose of the present study we limited participants to those who self-reported to be of European ethnicity. Exclusion criteria were: (1) a previous contact with specialist mental health services for psychotic symptoms outside of the study period at each site; (2) evidence of psychotic symptoms precipitated by an organic cause (ICD-10: F09); (3) transient psychotic symptoms resulting from acute intoxication (F1x.5); (4) severe learning disabilities, defined by an IQ less than 50 or diagnosis of intellectual disability (F70-F79); and (5) insufficient fluency of the primary language at each site to complete assessments.³¹

Ethical Approval

All participants who agreed to take part in the case-control study provided informed, written consent following full explanation of the study. Ethical approval for the study was provided by relevant local research ethics committees in each of the study sites.³²

Assessments

Socio-Demographic Characteristics. Using the Medical Research Council Sociodemographic modified Questionnaire version,³⁴ data on socio-demographic characteristics, including gender and country of birth, at the time of the first contact with mental health services for psychosis were collated at each research site. Age at first contact was defined as the age at which a patient was in contact with mental health services for the first time due to their psychotic symptoms. Ethnicity was self-ascribed from the 16 categories employed by the UK Census in 2001 (www.statistics.gov.uk/census 2001). Further educational attainment (no qualifications and school qualifications vs higher educational attachment which encompassed tertiary; vocational; undergraduate; postgraduate), employment status (unemployed vs employed [full- or part time] as a reference), living circumstances (currently living with people other than parents vs living alone or/and alone with children) and relationship status (ever vs never in a long-term [>1 year] relationship) were self-reported at first contact with services.

Clinical Measures. A modified version of the Nottingham Onset Schedule (NOS)³⁵ was used to measure DUP, based on the assessment interview and mental health records, and defined in weeks as the difference between the date of the first positive psychotic symptom (hallucination, delusion or thought disorder- rated as 4 [moderate-severe] or higher on the Positive and Negative Syndrome Scale [PANSS])³⁶ and the date of initiation of antipsychotic treatment.³⁷ The NOS scale provides a standardized and reliable way of recording early changes in psychosis and identifying relatively precise time points for measuring several durations in emerging psychosis.³⁵ The Operational Criteria Checklist (OPCRIT)³⁸ systems, whose reliability was assessed before and throughout the study ($k = 0.7$), was used by trained investigators to assess psychopathology in the first 4 weeks after the onset and generate research-based diagnoses based on ICD-10 diagnostic classification systems.³⁹ In the present study, diagnoses were grouped using ICD-10 codes into schizophrenia-spectrum disorders (F20-29), bipolar disorder (F30, F31), psychotic depression (F32, F33), and other psychosis.

Genetic Data

Samples were genotyped at the MRC Centre for Neuropsychiatric Genetics and Genomics in Cardiff (the United Kingdom) using a custom Illumina HumanCoreExome-24 BeadChip genotyping array covering 570038 genetic variants (Illumina Inc.).

Quality Control. Quality control (QC) entailed removing samples based on call rate (<0.99), genotype-phenotype mismatched information, suspected non-European ancestry, heterozygosity, and relatedness. Single-nucleotide polymorphism (SNPs) were

excluded if the minor allele frequency was 5%, if more than 2% of genotype data were missing and if the Hardy-Weinberg Equilibrium P -value $< 10^{-6}$; non-autosomal markers were also removed. The baseline characteristics of participants who were genotyped or were not genotyped are provided in [supplementary table 1](#). To account for any ancestry differences in genetic structures that could bias results, principal components analysis was conducted retaining top principal components (PCs).⁴⁰ Individuals of European ancestry were defined as having PC values within 6 SDs from the mean PC of the EUR in 1000G. Top 20 PCs were retained to adjust for possible population stratification in the association analyses.^{40,41}

Polygenic Scores. To calculate polygenic score for schizophrenia (SZ-PGS), bipolar disorder (BD-PGS), major depressive disorder (MDD-PGS) and intelligence (IQ-PGS), we used the summary statistics from the latest and largest genome-wide association studies^{10,22–24} utilizing PRSice⁴² where quality-controlled SNPs were pruned using clumping procedure which allowed to obtain SNPs in linkage equilibrium with an $r^2 < 0.25$ within a 250 kb window. Each PGS was calculated using subsets of the total SNPs based on the P -value threshold of .05. The selected P value threshold of .05 for SNP inclusion was chosen based on evidence showing that it explains the most variance.^{10,22–24} To aid interpretability of the results, all PGSs were standardized to a mean of 0 ($SD = 1$).

Statistical Analysis

All analyses reported in the present study were performed using RStudio version 4.0.3.⁴³

Imputation of Missing Values. In the present study, unemployed (22.9% missing), DUP (15.3% missing), diagnosis (1.8% missing) and living alone (1.0% missing) variable had missing values ([supplementary table 2](#)). To avoid using an unrepresentative sample of complete cases that may result in incorrect risk predictions,^{44,45} we conducted an imputation to handle the missing data. To impute the missing values, we employed missForest,⁴⁶ which is an iterative imputation method based on Random Forests. It handles continuous and categorical variables equally well and accommodates non-linear relation structures.⁴⁶ miss-Forest has been shown to outperform the well-known imputation methods, such as k -nearest neighbors and parametric multivariate imputation by chained equations.⁴⁶ To evaluate the quality of imputation, we estimated the imputation error Normalized Root Mean Squared Error (NRMSE) for continuous variables and proportion of falsely classified (PFC) for categorical variables.^{46,47} A value close to 0 represents an excellent performance, and a value of 1 indicates poor performance. The imputation of the missing values yielded a minimal error (NRMSE = 0.08%; PFC = 0.13%), highlighting that the imputed values were very closely aligned with

the observed values for both continuous and categorical variables. The distribution of the variables included in the analyses before and after the imputation are presented in [supplementary table 3](#).

Calculate Power and Predictive Accuracy of PGS. Using information on sample size (n), total number of independent markers in genotyping panel (m) and lower and upper P -values to select markers into polygenic score ($p0$, $p0.5$) we estimated the predictive accuracy (R^2) present in each PGS employed in the present study using Avengeme package implemented in R.⁴³ Consequently, using $n = 619$, and the number of SNPs included in PGS for schizophrenia ($m = 26\,281$), bipolar disorder ($m = 18\,092$), MDD ($m = 19\,508$) and intelligence ($m = 24\,386$), we estimated predictive accuracy for each PGS showing that SZ-PGS ($R^2 = 0.134$, $P = 7.40 \times 10^{-23}$), BD-PGS ($R^2 = 0.005$, $P = .044$), MDD-PGS ($R^2 = 0.036$, $P = 1.05 \times 10^{-6}$) and IQ-PGS ($R^2 = 0.077$, $P = 4.24 \times 10^{-13}$) had sufficient, as indicated by significant P -values, predictive accuracy to be employed in the analyses.

Regression Modeling. As the frequency distributions of DUP are severely skewed, DUP was normalized by taking the logarithm to base 10 (\log_{10} DUP) to allow the use of parametric regressions. Following log-transformation, \log_{10} DUP was normally distributed; distribution of DUP after normalization is presented in [supplementary figure 1](#); the results from the correlations between \log_{10} DUP and each PGS are provided in [supplementary table 5](#). For each PGS, 2 linear regression models were fitted to understand the role of covariates on the potential relationship of DUP with PGSs: Model 1: crude (unadjusted) model investigating an association between each PGS and DUP; Model 2: Model 1 plus adjusting for age at first contact with mental health services for psychosis, gender, genetic ancestry as measured with first 4 PCs, research sites and educational attainment. To measure prediction accuracy of each PGS, we utilized the incremental R^2 , which was calculated following the previously outlined steps.⁴⁸ Specifically, to calculate R^2 value for each model, we first regressed a phenotype on our set of controls without the PGS; we then re-ran the same regression but with the PGS included as a regressor.

Sensitivity Analyses. To examine whether our findings were applicable to individuals with a diagnosis of FEP disorders or were specific to people with a diagnosis of first-episode schizophrenia spectrum disorders only, we repeated the analyses limiting them to those who received the diagnosis of schizophrenia spectrum disorders on the first contact with mental health services. We further investigated if the results would remain the same using unimputed (complete cases) variables. As this was an exploratory study, which does not strictly require adjustment for multiple comparisons,⁴⁹ we did not employ correction for multiple testing. All tests for analyses were 2-tailed; P -values $\leq .05$ were considered statistically significant.

Results

Sample Characteristics

The demographic characteristics of the analytic sample of FEP cases are presented in [table 1](#). The sample comprised 619 (86.6% of $N = 715$) individuals of European ancestry for whom quality-controlled genome-wide genotyping and DUP were available. Those participants who were included in the study or excluded from the final cohort did not differ in terms of DUP, gender, marital status, employment, living arrangement, and diagnoses; however, the former group included participants who were younger ($t_{(1112.5)} = -2.31, P = .021$) and had a lower educational attainment ($\chi^2_{(1)} = 4.72, P = .030$) compared to those who were included in the study ([supplementary table 4](#)). The median age at first contact was 29.0 years (IQR = 22.0–38.0). Of the entire sample, 37.3% ($N = 227/669$) had diagnoses of first-episode schizophrenia spectrum disorders, 63.6% ($N = 394$) were men, 37.3% ($N = 178$) were unemployed and 18.8% lived alone at the time of the first contact with mental health services for psychosis.

Length of DUP by European Countries and FEP Diagnoses

The median length of DUP in the whole sample was 10.0 weeks (interquartile range [IQR] = 3.8–30.8). DUP did not differ by countries: France (median = 12.3 weeks, IQR = 80.8), the United Kingdom (median = 10.9 weeks, IQR = 2.42–51.93), The Netherlands (median = 10.3 weeks, IQR = 3.42–29.63), Italy (median = 8.1 weeks,

Table 1. Baseline Sociodemographic and Clinical Characteristics of First-Presentation Psychosis Patients

Baseline Sample Characteristics	Total Sample ($N = 619$)
	N (%) / mean (SD)
Age (y)	31.5 (10.9)
DUP (wk)	62.5 (191.6)
Male gender	394 (63.6)
Not married	444 (72.1)
Unemployed	178 (37.3)
Living alone	115 (18.8)
Low educational attainment	88 (14.3)
Diagnosis	
Schizophrenia spectrum	250 (11.0)
Bipolar Disorder	67 (11.0)
Psychotic depression	74 (12.2)
Other psychosis	217 (35.7)
Country of data collection	
The United Kingdom	99 (16.0)
Holland	133 (21.5)
Spain	151 (24.4)
France	24 (6.8)
Italy	103 (16.6)
Brazil	91 (14.7)

Note: DUP, duration of untreated psychosis.

IQR = 3.85–18.80), Spain (median = 8.7 weeks, IQR = 4.27–34.83), and Brazil (median = 9.4 weeks, IQR = 4.13–24.08) (Kruskal-Wallis₍₆₎ = 4.61, $P = .595$). When DUP was stratified by diagnoses, the longest DUP was observed in patients with diagnosis of schizophrenia spectrum (median = 20.8 weeks, IQR = 10.11–80.07) followed by psychotic depression (median = 11.5 weeks, IQR = 6.41–22.65) and bipolar disorders (median = 4.42 weeks, IQR = 1.71–8.69) (Kruskal-Wallis₍₃₎ = 110.0, $P = 1.09 \times 10^{-23}$).

Associations Between PGSs and DUP

Associations between length of DUP and PGS in patients with FEP are presented in [table 2](#). One SD increase in SZ-PGS was not significantly associated with the length of DUP in participants with a diagnosis of FEP disorders (Model 2: $\beta_{\text{adjusted}} = -0.11$, 95% CI = -0.341 – -0.131 , $R^2 = 0.023$). Similarly, there were no statistically significant associations between DUP and BD-PGS (Model 2: $\beta_{\text{adjusted}} = 0.050$, 95% CI = -0.123 – -0.223 , $R^2 = 0.022$), MDD-PGS (Model 2: $\beta_{\text{adjusted}} = 0.036$, 95% CI = -0.094 – -0.167 , $R^2 = 0.022$) or IQ-PGS (Model 2: $\beta_{\text{adjusted}} = -0.017$, 95% CI = -0.160 – -0.125 , $R^2 = 0.022$) in participants with a diagnosis of FEP disorders. These results did not differ from those observed in Model 1 (ie, the unadjusted model) and when complete-case (unimputed) data were employed to run the models ([supplementary table 6](#)). When analyses were limited to participants with a diagnosis of first-episode schizophrenia spectrum, we did not find significant associations between each polygenic score and DUP in unadjusted and fully adjusted models ([supplementary table 7](#)).

Discussion

To our knowledge, this is the first study investigating the relationship of polygenic propensity for schizophrenia, bipolar disorder, major depressive disorder, and intelligence with duration of untreated psychosis.

Consistent with previous reports,^{50–52} our findings showed that individuals with a diagnosis of FEP disorders had to endure a prolonged period coping with symptoms of psychosis without seeking appropriate treatments; though, this was heavily skewed with a smaller subset of participants experiencing over a year before first contact with mental health services. Similar to previous reports,^{17,30} we observed that the median length of DUP in participants with schizophrenia spectrum disorders was significantly longer when compared to all other psychoses. These observed delays highlight that there is still a great need to improve recognition of the symptoms of FEP, including schizophrenia and pathways to care.

The neurodevelopmental theory of schizophrenia posits that genetic factors interfere with early brain development leading to the development of schizophrenia

Table 2. Associations Between Length of Untreated Psychosis and PGS in Patients With FEP

PGS	Model 1			Model 2		
	β (95% CI)	P-Value	Model Fit	β (95% CI)	P-Value	Model Fit
SZ-PGS	0.052 (-0.089–0.194)	.468	$R^2 = 0.001$	-0.110 (-0.341–0.131)	.467	$R^2 = 0.023$
BD-PGS	-0.020 (-0.161–0.122)	.785	$R^2 = 0.000$	0.050 (-0.123–0.223)	.389	$R^2 = 0.022$
MDD-PGS	0.060 (-0.081–0.201)	.405	$R^2 = 0.001$	0.036 (-0.094–0.167)	.149	$R^2 = 0.022$
IQ-PGS	0.008 (-0.133–0.150)	.907	$R^2 = 0.000$	-0.017 (-0.160–0.125)	.776	$R^2 = 0.022$

Note: Effect size is indicated by β coefficient from the linear regression model; the presented β coefficient is standardized. CI, confidence interval; SZ-PGS, polygenic score for schizophrenia; BD-PGS, bipolar disorders; MDD-PGS, major depressive disorder; IQ-PGS, intelligence; PGS, polygenic scores; FEP, first episode psychosis. Model 1: crude (unadjusted) model investigating an association between each PGS and DUP; Model 2: Model 1 plus adjusting for age at first contact with mental health services for psychosis, gender, genetic ancestry as measured with first 4 principal components, research site and educational attainment.

symptoms.⁵³ This, in combination of accumulated evidence for polygenicity of schizophrenia,¹⁰ alluded to a possibility that the length of DUP might also be influenced by additive effect of multiple common genetic markers linked to schizophrenia.⁸ However, this hypothesis was not confirmed by our findings. We further considered that high genetic predisposition for either bipolar disorder or major depression disorder might be associated with DUP in people with a diagnosis of FEP disorders. Once again, our findings were negative, as they were for PGS for intelligence. It may be argued that a limited power might have led to these non-significant results. Because the PGS employed in this study were built using the results from most recent and largest GWAS meta-analyses, our analyses were not constrained by our sample size.^{43,54} Nonetheless, to ensure we captured the true polygenic contribution to DUP, we undertook calculations of power for each polygenic score, which revealed that there was considerable predicative power in each PGS to detect potential associations. Our results are further in line with a recent literature review highlighting that evidence of an association between DUP and brain structure in people with a diagnosis FEP disorders was minimal.⁵⁵ It is further argued that any relationships observed between untreated psychosis and psychosis illness course appears to be explained by lead-time bias.⁷ Accordingly, those with a short DUP are in an earlier stage and therefore are likely to have better outcomes than those with a long DUP, who are in a later stage.⁷ Cumulatively, our findings shed some doubt on the notion that genetic variation has substantial impact on DUP.⁵⁶

In light of these findings, a discussion of some alternative theories explaining the length of DUP is warranted. It has been suggested that the length of delay from first manifestation of psychotic symptoms to initiation of adequate treatment may be influenced by factors related to the organization of mental health system and process of referral to an appropriate service FEP. Reduced allocated resources for early intervention services⁵⁷ and limited availability of care⁵⁸ may also be important contributing factors to longer DUP. The lack of knowledge of what

constitutes psychosis onset^{5,37} and what help may be available for people affected by early psychosis and their families⁵⁹ were shown to be important factors influencing DUP.^{5,37} The longer delays to seeking help for first-episode psychotic disorders were further linked fear of stigma. Therefore, DUP may be significantly reduced through educational and anti-stigmatizing campaign about the signs of early psychosis targeted at health care providers, public, and schools increasing the motivation to seek treatment.⁵⁸ Although evidences regarding successfulness of specific interventions in reducing DUP are still lacking and largely non replicated,⁶⁰ our findings should encourage the identification of potentially effective initiatives.

Methodological Strengths and Limitations

This is an extensive multi-site study of FEP with comprehensive data on a variety of environmental and genetic factors. The study included all incidence cases from well-defined catchment areas in 17 sites across 6 countries. As our analyses were focused on people with a diagnosis of FEP, the findings reported in the present study are less likely to be biased toward patients who experience multiple hospital admissions.^{61,62} Given that our study was carried out in major urban and rural sites with heterogeneous populations suggest that the generalizability of our findings may extend to other centers with similar population profiles. Finally, because the calculation of PGS is based on well-powered GWASs, we did not require a large sample to test our hypotheses, which was further confirmed by estimated predictive power in each PGS employed in the analyses.

Nonetheless, important methodological considerations warrant a discussion. While it is likely most individuals who develop a psychotic disorder do present to services, at least in sites with well-developed public health systems, some who do not present will be missed and this may introduce selection biases.³¹ Variations in referral procedures of patients with psychosis from primary to secondary mental health care settings and in the organization of secondary mental health care services across

catchment areas may have influenced the identification of cases.³¹ Even though robust imputation methods have been employed to deal with missing values, the percentage of missingness in our variables, though lower than previously, is a notable issue in the present study. Although the median age of our sample was consistent with that of other very large samples of individuals with FEP collated in Europe, Australia and some studies from North American,⁶³⁻⁶⁵ it may still be higher compared to other studies from the US. Thus, we urge caution when generalizing our findings to all patients with FEP across the continents. It may be argued that the length of DUP observed in our study was shorter than that reported in some other studies,⁵⁰⁻⁵² which in turn may have reduced the likelihood of finding a significant association with PGS. The poor generalizability of genetic studies across populations is also noteworthy.³³ This is because the construction of PGSs is largely dependent on the availability of the summary statistics from genome-wide association studies (GWASs). However, around 79% of all GWAS participants are of European descent despite making up only 16% of the global population.³³ Given genetic risk is different in European and non-European individuals, further work is necessary to develop PGSs model in non-White populations. Because our analyses were restricted to individuals of European ethnicity, our results do not shed light on the associations of genetic predisposition to the major psychiatric conditions and DUP among people of non-European ethnicity. This is an important limitation as Black-African and Black-Caribbean groups were shown to have a significantly different length of DUP relative to White groups,⁶⁶ which in turn may reflect differences in pathways to care experienced by some ethnic minority groups.⁶⁷ Finally, by their design, PGSs do not capture other structural variants beyond common genetic markers of relatively small effects, such as rare variants, poorly tagged or multiple independent variants, gene-by-gene interaction and gene-by-environment interplay.⁶⁸

Conclusion

Although our findings are specific to individuals from European populations, our results suggest there are not strong genetic risk factors underlying duration of untreated psychosis, underscoring the importance of effective educational efforts directed towards the public, the schools, and the health professionals about first onset of psychotic disorders.

Supplementary Material

Supplementary material is available at *Schizophrenia Bulletin*.

EU-GEI Collaborators: Hubbard Kathryn¹, Beards Stephanie¹, Stilo Simona A.², Parellada Mara³, Cuadrado Pedro⁴, Solano José Juan Rodríguez⁵,

David Fraguas⁵, Andreu-Bernabeu Álvaro⁵, Carracedo Angel⁶, Bernardo Enrique García⁷, Roldán Laura³, López Gonzalo³, Cabrera Bibiana⁸, Nacher Juan⁹, Garcia-Portilla Paz¹⁰, Costas Javier⁶, Jiménez-López Estela¹¹, Matteis Mario³, Castro Marta Rapado³, González Emiliano³, Martínez Covadonga³, Sánchez Emilio⁷, Durán-Cutilla Manuel⁷, Franke Nathalie¹², Termorshuizen Fabian¹³,¹⁴, van Dam Daniella¹², van der Ven Elsje^{13,14}, Messchaert Elles¹⁴, Leboyer Marion^{15,16,17,18}, Schürhoff Franck^{15,16,17,18}, Jamain Stéphane^{16,17,18}, Baudin Grégoire^{15,16}, Ferchiou Aziz^{15,16}, Pignon Baptiste^{15,16,18}, Richard Jean- Romain^{16,18}, Charpeaud Thomas^{18,19,21}, Tronche Anne-Marie^{18,19,21}, Frijda Flora²², La Barbera Daniele^{22,23}, Marrazzo Giovanna²³, Sideli Lucia²², Sartorio Crocettarachele^{22,23}, Ferraro Laura²², Seminerio Fabio²², Loureiro Camila Marcelino^{24,25}, Shuhama Rosana^{24,25}, Ruggeri Mirella²⁶, LaSalvia Antonio²⁶, Bonetto Chiara²⁶, and Cristofalo Doriana²⁶. Corresponding affiliations: ¹Department of Health Service and Population Research, Institute of Psychiatry, King's College London, De Crespigny Park, Denmark Hill, London, United Kingdom, SE5 8AF; ²Department of Psychosis Studies, Institute of Psychiatry, King's College London, De Crespigny Park, Denmark Hill, London, United Kingdom SE5 8AF; ³Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM (CIBERSAM), C/Doctor Esquerdo 46, 28007 Madrid, Spain; ⁴Villa de Vallecas Mental Health Department, Villa de Vallecas Mental Health Centre, Hospital Universitario Infanta Leonor / Hospital Virgen de la Torre, C/San Claudio 154, 28038 Madrid, Spain; ⁵Puente de Vallecas Mental Health Department, Hospital Universitario Infanta Leonor / Hospital Virgen de la Torre, Centro de Salud Mental Puente de Vallecas, C/Peña Gorbea 4, 28018 Madrid, Spain; ⁶Fundación Pública Galega de Medicina Xenómica, Hospital Clínico Universitario, Choupana s/n, 15782 Santiago de Compostela, Spain; ⁷Department of Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM (CIBERSAM), C/Doctor Esquerdo 46, 28007 Madrid, Spain; ⁸Department of Psychiatry, Hospital Clinic, IDIBAPS, Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Universidad de Barcelona, C/Villarroel 170, escalera 9, planta 6, 08036 Barcelona, Spain; ⁹Neurobiology Unit, Program in Neurosciences and Interdisciplinary Research Structure for Biotechnology and Biomedicine (BIOTECMED), Universitat de València, Burjassot, Spain Biomedical Research Networking Centre in Mental Health (CIBERSAM), Madrid, Spain. Biomedical Research Institute INCLIVA, Valencia, Spain; ¹⁰Department of Medicine, Psychiatry

Area, School of Medicine, Universidad de Oviedo, Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), C/Julián Clavería s/n, 33006 Oviedo, Spain; ¹¹Department of Psychiatry, Servicio de Psiquiatría Hospital “Virgen de la Luz,” C/Hermanidad de Donantes de Sangre, 16002 Cuenca, Spain; ¹²Department of Psychiatry, Early Psychosis Section, Academic Medical Centre, University of Amsterdam, Meibergdreef 5, 1105 AZ Amsterdam, The Netherlands; ¹³Rivierduinen Centre for Mental Health, Leiden, Sandiforddreef 19, 2333 ZZ Leiden, The Netherlands; ¹⁴Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, South Limburg Mental Health Research and Teaching Network, Maastricht University Medical Centre, P.O. Box 616, 6200 MD Maastricht, The Netherlands; ¹⁵AP-HP, Groupe Hospitalier “Mondor,” Pôle de Psychiatrie, 51 Avenue de Maréchal de Lattre de Tassigny, 94010 Créteil, France; ¹⁶INSERM, U955, Equipe 15, 51 Avenue de Maréchal de Lattre de Tassigny, 94010 Créteil, France; ¹⁷Faculté de Médecine, Université Paris-Est, 51 Avenue de Maréchal de Lattre de Tassigny, 94010 Créteil, France; ¹⁸Fondation Fondamental, 40 Rue de Mesly, 94000 Créteil, France; ¹⁹CMP B CHU, BP 69, 63003 Clermont Ferrand, Cedex 1, France; ²⁰EPS Maison Blanche, Paris 75020, France; ²¹Université Clermont Auvergne, EA 7280, Clermont-Ferrand, 63000, France; ²²Department of Experimental Biomedicine and Clinical Neuroscience, Section of Psychiatry, University of Palermo, Via G. La Loggia n.1, 90129 Palermo, Italy; ²³Unit of Psychiatry, “P. Giaccone” General Hospital, Via G. La Loggia n.1, 90129 Palermo, Italy; ²⁴Departamento de Neurociências e Ciências do Comportamento, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Av. Bandeirantes, 3900 - Monte Alegre-CEP 14049-900, Ribeirão Preto, SP, Brasil; ²⁵Núcleo de Pesquisa em Saúde Mental Populacional, Universidade de São Paulo, Avenida Doutor Arnaldo 455, CEP 01246-903, SP, Brasil; ²⁶Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement, University of Verona, Piazzale L.A. Scuro 10, 37134 Verona, Italy.

Acknowledgments

R.M.M. has received honoraria from Janssen, Sunovian, Lundbeck and Otsuka. M.B. has been a consultant for, received grant/research support and honoraria from, and been on the speakers/advisory board of ABBiotics, Adamed, Angelini, Casen Recordati, Janssen-Cilag, Menarini, Rovi and Takeda. Other authors declare that they have no conflict of interest. All other authors declare no conflict of interest.

Funding

O.A. is funded by the National Institute for Health Research (NIHR) (NIHR Post-Doctoral Fellowship

- PDF-2018-11-ST2-020). V.R. was funded by a PhD scholarship supported by Lord Leverhulme’s Charitable Trust and by the Velvet Foundation. M.D.F. is funded by Clinician Scientist Medical Research Council fellowship (project reference MR/M008436/1). D.Q. is funded by Post-Doctoral Guarantors of Brain Clinical Fellowship. The EU-GEI Project is funded by the European Community’s Seventh Framework Programme under grant agreement No. HEALTH-F2-2010-241909 (Project EU-GEI). The Brazilian study was funded by the São Paulo Research Foundation under grant number 2012/0417-0. J.B.K. was supported by National Institute for Health Research, University College London Hospital, Biomedical Research Centre. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health and Social Care. Funders were not involved in the design and conduct of the study; collection, management, analysis and interpretation of the data; preparation, review or approval of the manuscript and decision to submit the manuscript for publication. The funding organizations had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

References

1. Zipursky RB, Reilly TJ, Murray RM. The myth of schizophrenia as a progressive brain disease. *Schizophr Bull.* 2013;39(6):1363–1372.
2. Perkins DO, Gu H, Boteva K, Lieberman JA. Relationship between duration of untreated psychosis and outcome in first-episode schizophrenia: a critical review and meta-analysis. *Am J Psychiatry.* 2005;162(10):1785–1804.
3. Lieberman JA, Perkins D, Belger A, et al. The early stages of schizophrenia: speculations on pathogenesis, pathophysiology, and therapeutic approaches. *Biol Psychiatry.* 2001;50(11):884–897.
4. Marshall M, Lewis S, Lockwood A, Drake R, Jones P, Croudace T. Association between duration of untreated psychosis and outcome in cohorts of first-episode patients: a systematic review. *Arch Gen Psychiatry.* 2005;62(9):975–983.
5. McGlashan TH. Duration of untreated psychosis in first-episode schizophrenia: marker or determinant of course? *Biol Psychiatry.* 1999;46(7):899–907.
6. Lieberman JA, Fenton WS. Delayed detection of psychosis: causes, consequences, and effect on public health. *Am J Psychiatry.* 2000;157(11):1727–1730.
7. Jonas KG, Fochtmann LJ, Perlman G, et al. Lead-time bias confounds association between duration of untreated psychosis and illness course in schizophrenia. *Am J Psychiatry.* 2020;177(4):327–334.
8. Verdoux H, Liraud F, Bergey C, Assens F, Abalan F, van Os J. Is the association between duration of untreated psychosis and outcome confounded? A two year follow-up study of first-admitted patients. *Schizophr Res.* 2001;49(3):231–241.

9. Harrigan SM, McGorry PD, Krstev H. Does treatment delay in first-episode psychosis really matter? *Psychol Med*. 2003;33(1):97–110.
10. Schulze TG, Akula N, Breuer R, et al.; Bipolar Genome Study. Molecular genetic overlap in bipolar disorder, schizophrenia, and major depressive disorder. *World J Biol Psychiatry*. 2014;15(3):200–208.
11. Schizophrenia Working Group of the Psychiatric Genomics Consortium, Ripke S, James TRW, O'Donovan MC. Mapping genomic loci prioritises genes and implicates synaptic biology in schizophrenia. *medRxiv The preprint server for health science*. 2020. doi:10.1101/2020.09.12.20192922
12. Wray NR, Lee SH, Mehta D, Vinkhuyzen AA, Dudbridge F, Middeldorp CM. Research review: polygenic methods and their application to psychiatric traits. *J Child Psychol Psychiatry*. 2014;55(10):1068–1087.
13. Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF, Sklar P. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*. 2009;460(7256):748–752.
14. Dudbridge F. Power and predictive accuracy of polygenic risk scores. *PLoS Genet*. 2013;9(3):e1003348.
15. Fanous AH, Zhou B, Aggen SH, et al.; Schizophrenia Psychiatric Genome-Wide Association Study (GWAS) Consortium. Genome-wide association study of clinical dimensions of schizophrenia: polygenic effect on disorganized symptoms. *Am J Psychiatry*. 2012;169(12):1309–1317.
16. Howes OD, Whitehurst T, Shatalina E, et al. The clinical significance of duration of untreated psychosis: an umbrella review and random-effects meta-analysis. *World Psychiatry*. 2021;20(1):75–95.
17. Bechard-Evans L, Schmitz N, Abadi S, Joobar R, King S, Malla A. Determinants of help-seeking and system related components of delay in the treatment of first-episode psychosis. *Schizophr Res*. 2007;96(1-3):206–214.
18. Aleman A, Agrawal N, Morgan KD, David AS. Insight in psychosis and neuropsychological function: meta-analysis. *Br J Psychiatry*. 2006;189:204–212.
19. Romm KL, Rossberg JI, Berg AO, et al. Depression and depressive symptoms in first episode psychosis. *J Nerv Ment Dis*. 2010;198(1):67–71.
20. Oyffe I, Shwizer R, Stolovy T. The association between diagnosis, treatment delay and outcome among patients with bipolar disorders. *Psychiatr Q*. 2015;86(1):95–105.
21. Angst J, Gamma A, Lewinsohn P. The evolving epidemiology of bipolar disorder. *World Psychiatry*. 2002;1(3):146–148.
22. Savage JE, Jansen PR, Stringer S, et al. Genome-wide association meta-analysis in 269,867 individuals identifies new genetic and functional links to intelligence. *Nat Genet*. 2018;50(7):912–919.
23. Stahl EA, Breen G, Forstner AJ, et al. Genome-wide association study identifies 30 loci associated with bipolar disorder. *Nat Genet*. 2019;51(5):793–803.
24. Wray NR, Ripke S, Mattheisen M, et al.; eQTLGen; 23andMe; Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium. Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet*. 2018;50(5):668–681.
25. Lichtenstein P, Yip BH, Björk C, et al. Common genetic determinants of schizophrenia and bipolar disorder in Swedish families: a population-based study. *Lancet*. 2009;373(9659):234–239.
26. Wray NR, Yang J, Hayes BJ, Price AL, Goddard ME, Visscher PM. Pitfalls of predicting complex traits from SNPs. *Nat Rev Genet*. 2013;14(7):507–515.
27. Benjamin DJ, Cesarini D, Chabris CF, et al. The promises and pitfalls of geneoconomics*. *Annu Rev Econom*. 2012;4:627–662.
28. Belsky DW, Harden KP. Phenotypic annotation: using polygenic scores to translate discoveries from genome-wide association studies from the top down. *Curr Dir Psychol Sci*. 2019;28:82–90.
29. Freese J. The arrival of social science genomics. *Contemp Sociol*. 2018;47(5):524–536.
30. Schimmelmann BG, Huber CG, Lambert M, Cotton S, McGorry PD, Conus P. Impact of duration of untreated psychosis on pre-treatment, baseline, and outcome characteristics in an epidemiological first-episode psychosis cohort. *J Psychiatr Res*. 2008;42(12):982–990.
31. Gayer-Anderson C, Jongsma HE, Di Forti M, et al.; EU-GEI WP2 Group. The European Network of National Schizophrenia Networks Studying Gene-Environment Interactions (EU-GEI): incidence and first-episode case-control programme. *Soc Psychiatry Psychiatr Epidemiol*. 2020;55(5):645–657.
32. Jongsma HE, Gayer-Anderson C, Lasalvia A, et al.; European Network of National Schizophrenia Networks Studying Gene-Environment Interactions Work Package 2 (EU-GEI WP2) Group. Treated incidence of psychotic disorders in the multinational EU-GEI study. *JAMA Psychiatry*. 2018;75(1):36–46.
33. Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Clinical use of current polygenic risk scores may exacerbate health disparities. *Nat Genet*. 2019;51(4):584–591.
34. Mallett R, Leff J, Bhugra D, Pang D, Zhao JH. Social environment, ethnicity and schizophrenia. A case-control study. *Soc Psychiatry Psychiatr Epidemiol*. 2002;37(7):329–335.
35. Singh SP, Cooper JE, Fisher HL, et al. Determining the chronology and components of psychosis onset: the Nottingham Onset Schedule (NOS). *Schizophr Res*. 2005;80(1):117–130.
36. Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull*. 1987;13(2):261–276.
37. Morgan C, Abdul-Al R, Lappin JM, et al.; AESOP Study Group. Clinical and social determinants of duration of untreated psychosis in the AESOP first-episode psychosis study. *Br J Psychiatry*. 2006;189:446–452.
38. McGuffin P, Farmer A, Harvey I. A polydiagnostic application of operational criteria in studies of psychotic illness. Development and reliability of the OPCRIT system. *Arch Gen Psychiatry*. 1991;48(8):764–770.
39. Quattrone D, Di Forti M, Gayer-Anderson C, et al.; EU-GEI WP2 Group. Transdiagnostic dimensions of psychopathology at first episode psychosis: findings from the multinational EU-GEI study. *Psychol Med*. 2019;49(8):1378–1391.
40. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet*. 2006;38(8):904–909.
41. Wang D, Sun Y, Stang P, Berlin JA, Wilcox MA, Li Q. Comparison of methods for correcting population stratification in a genome-wide association study of rheumatoid arthritis: principal-component analysis versus multidimensional scaling. *BMC Proc*. 2009;3(Suppl 7):S109.

42. Euesden J, Lewis CM, O'Reilly PF. PRSice: Polygenic Risk Score software. *Bioinformatics*. 2015;31(9):1466–1468.
43. RStudioTeam. Boston, MA: RStudio: Integrated Development for R. Inc; 2016. <http://www.rstudio.com/>.
44. Moons KG, Donders RA, Stijnen T, Harrell FE Jr. Using the outcome for imputation of missing predictor values was preferred. *J Clin Epidemiol*. 2006;59(10):1092–1101.
45. Zhao Y, Long Q. Multiple imputation in the presence of high-dimensional data. *Stat Methods Med Res*. 2016;25(5):2021–2035.
46. Stekhoven DJ, Bühlmann P. MissForest—non-parametric missing value imputation for mixed-type data. *Bioinformatics*. 2012;28(1):112–118.
47. Oba S, Sato MA, Takemasa I, Monden M, Matsubara K, Ishii S. A Bayesian missing value estimation method for gene expression profile data. *Bioinformatics*. 2003;19(16):2088–2096.
48. Lee JJ, Wedow R, Okbay A, et al.; 23andMe Research Team; COGENT (Cognitive Genomics Consortium); Social Science Genetic Association Consortium. Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat Genet*. 2018;50(8):1112–1121.
49. Bender R, Lange S. Adjusting for multiple testing—when and how? *J Clin Epidemiol*. 2001;54(4):343–349.
50. Barnes TR, Hutton SB, Chapman MJ, Mutsatsa S, Puri BK, Joyce EM. West London first-episode study of schizophrenia. Clinical correlates of duration of untreated psychosis. *Br J Psychiatry*. 2000;177:207–211.
51. Loebel AD, Lieberman JA, Alvir JM, Mayerhoff DI, Geisler SH, Szymanski SR. Duration of psychosis and outcome in first-episode schizophrenia. *Am J Psychiatry*. 1992;149(9):1183–1188.
52. Hoff AL, Sakuma M, Razi K, Heydebrand G, Csernansky JG, DeLisi LE. Lack of association between duration of untreated illness and severity of cognitive and structural brain deficits at the first episode of schizophrenia. *Am J Psychiatry*. 2000;157(11):1824–1828.
53. Murray RM, Lewis SW. Is schizophrenia a neurodevelopmental disorder? *Br Med J (Clin Res Ed)*. 1987;295(6600):681–682.
54. Choi SW, Mak TS, O'Reilly PF. Tutorial: a guide to performing polygenic risk score analyses. *Nat Protoc*. 2020;15(9):2759–2772.
55. Anderson KK, Rodrigues M, Mann K, et al. Minimal evidence that untreated psychosis damages brain structures: a systematic review. *Schizophr Res*. 2015;162(1-3):222–233.
56. Kraguljac NV, Anthony T, Morgan CJ, Jindal RD, Burger MS, Lahti AC. White matter integrity, duration of untreated psychosis, and antipsychotic treatment response in medication-naïve first-episode psychosis patients. *Mol Psychiatry*. 2020. doi:10.1038/s41380-020-0765-x
57. O'Donoghue B, Lyne JP, Renwick L, et al. Neighbourhood characteristics and the incidence of first-episode psychosis and duration of untreated psychosis. *Psychol Med*. 2016;46(7):1367–1378.
58. Johannessen JO, McGlashan TH, Larsen TK, et al. Early detection strategies for untreated first-episode psychosis. *Schizophr Res*. 2001;51(1):39–46.
59. Larsen TK, Johannessen JO, Opjordsmoen S. First-episode schizophrenia with long duration of untreated psychosis. Pathways to care. *Br J Psychiatry Suppl*. 1998;172(33):45–52.
60. Oliver D, Davies C, Crossland G, et al. Can we reduce the duration of untreated psychosis? A systematic review and meta-analysis of controlled interventional studies. *Schizophr Bull*. 2018;44(6):1362–1372.
61. Ajnakina O, Stubbs B, Francis E, et al. Hospitalisation and length of hospital stay following first-episode psychosis: systematic review and meta-analysis of longitudinal studies. *Psychol Med*. 2020;50(6):991–1001.
62. Lally J, Ajnakina O, Stubbs B, et al. Remission and recovery from first-episode psychosis in adults: systematic review and meta-analysis of long-term outcome studies. *Br J Psychiatry*. 2017;211(6):350–358.
63. Morgan C, Lappin J, Heslin M, et al. Reappraising the long-term course and outcome of psychotic disorders: the AESOP-10 study. *Psychol Med*. 2014;44(13):2713–2726.
64. Ajnakina O, Lally J, Di Forti M, et al. Patterns of illness and care over the 5 years following onset of psychosis in different ethnic groups; the GAP-5 study. *Soc Psychiatry Psychiatr Epidemiol*. 2017;52(9):1101–1111.
65. Norman RM, Manchanda R, Windell D. The prognostic significance of early remission of positive symptoms in first treated psychosis. *Psychiatry Res*. 2014;218(1-2):44–47.
66. Schoer N, Huang CW, Anderson KK. Differences in duration of untreated psychosis for racial and ethnic minority groups with first-episode psychosis: an updated systematic review and meta-analysis. *Soc Psychiatry Psychiatr Epidemiol*. 2019;54(10):1295–1298.
67. Anderson KK, Flora N, Archie S, Morgan C, McKenzie K. A meta-analysis of ethnic differences in pathways to care at the first episode of psychosis. *Acta Psychiatr Scand*. 2014;130(4):257–268.
68. Reynolds CA, Finkel D. A meta-analysis of heritability of cognitive aging: minding the “missing heritability” gap. *Neuropsychol Rev*. 2015;25(1):97–112.