

# Silylium-Catalyzed Regio- and Stereoselective Carbosilylation of Ynamides with Allylic Trimethylsilanes

Paz Yepes, Ángel L. Suárez-Sobrino, Miguel A. Rodríguez, and Alfredo Ballesteros\*



Cite This: *Org. Lett.* 2023, 25, 1020–1024



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

**ABSTRACT:** The regio- and stereoselective carbosilylation of tosylynamides with allylic trimethylsilanes takes place under mild conditions in the presence of catalytic TMSNTf<sub>2</sub> or HNTf<sub>2</sub> to give (*Z*)- $\alpha$ -allyl- $\beta$ -trimethylsilylenamides with good yields. Theoretical calculations show the activation of the C–C triple bond of the ynamides by the trimethylsilylium ion and formation of a  $\beta$ -trimethylsilylketenimonium cation. Further transformations of the products demonstrate the synthetic utility of this reaction.




Silylium ion has been consolidated in the past two decades as a potent catalyst in organic synthesis.<sup>1</sup> Its strong Lewis acid character is reflected in its high affinity not only to  $\sigma$ - but also to  $\pi$ -bases. This makes silylium ion a simpler and more sustainable alternative to catalytic metal salts or transition metal complexes for the activation of C–C multiple bonds. Since the pioneering work of Lambert et al., employing 1,1-disubstituted alkenes,<sup>2</sup> several examples of silylium-catalyzed hydro-<sup>3–6</sup> and carbosilylation<sup>7</sup> of C–C double bonds have appeared. However, there are few precedents related to the activation of triple C–C bonds with silylium ion.<sup>8</sup> In this context, Kawashima et al. recently described a silylium catalyzed intermolecular silylation of an arylalkyne<sup>9</sup> to form a  $\beta$ -silyl stabilized vinylcation, that was subsequently intercepted by an intramolecular Friedel–Crafts ring closure.


Exploring new ways for the activation of electron rich triple C–C bonds, ynamides could be good candidates to broad the scope of silylium ion catalysis. The tendency of ynamides to be activated by electrophilic species such as acids and transition metals,<sup>10</sup> and the polarization of their C–C triple bond allow many regioselective reactions.<sup>10,11</sup> Moreover, the 1,2 functionalization of ynamides offers the possibility to obtain functionalized and highly substituted nitrogenated alkenes.<sup>12</sup> Thus, the silylation of these compounds represents an entry to nitrogen-substituted vinylsilanes,<sup>13</sup> of great value in organic synthesis.

Several methods to install a silyl group in one of the carbon atoms of ynamides have been described (Figure 1). Thus,  $\alpha,\beta$ -silylmetalation and subsequent attack of an electrophile to the metal position is the most common method for this purpose (Figure 1A); for instance, the silylcupration<sup>14</sup> and the Pd-catalyzed silylstannation<sup>15</sup> of ynamides result in  $\alpha$ -metalated (*Z*)- $\beta$ -silylenamides; alternatively, Pd-catalyzed silylboration leads to  $\beta$ -metalated (*Z*)- $\alpha$ -silylenamides;<sup>16</sup> finally,  $\alpha$ -metalated (*E*)- $\beta$ -silylenamides can be obtained from a *trans*-selective radical silylzincation of ynamides.<sup>17</sup>


## A. Silylmetalation (ref 14–17)



## B. Hydrosilylation (ref 18,19)



## C. Silylcyanation (ref 20)



## D. Allylsilylation (this work)





Figure 1. Ynamide silylation methods.

Apart from the silylmetalation approach,  $\beta$ -silyl-(*Z*)-enamides can also selectively be obtained by hydrosilylation of ynamides using a rhodium complex<sup>18</sup> or tris-(pentafluorophenyl)borane<sup>19</sup> as catalyst; in both cases, the hydride abstraction from the silane by the corresponding catalyst leads to a silylium ion, responsible for the formation of a  $\beta$ -silyl ketenimonium intermediate I (Figure 1B). Very

Received: January 20, 2023

Published: February 7, 2023




recently, a palladium catalyzed silylcyanation was described; the reaction proceeds in a stereo- and regioselective way through a  $\beta$ -palladium enamide intermediate **II** (Figure 1C).<sup>20</sup>

On the other hand, the precedented reactions of allylsilanes with C–C multiple bonds catalyzed by Brønsted or Lewis acids<sup>21–23</sup> and the possibility of self-regeneration of catalytic silylium moved us to choose allylsilane derivatives as carbon nucleophile counterparts for our study on the catalytic carbosilylation of ynamides (Figure 1D). Therefore, herein we develop a regio- and stereoselective allylsilylation of ynamides, using catalytic silylium ion, an alternative to other species such as metal salts and transition metal complexes.

Our first experiments focused on the reaction between tosylamide **1a** and allyltrimethylsilane **2a** in 1,2-dichloroethane (DCE) using a direct silylium ion freshly prepared source like *N*-trimethylsilyl bis(trifluoromethanesulfonyl)imide, TMSNTf<sub>2</sub>,<sup>24</sup> or an acid like bis(trifluoromethanesulfonyl)imide, HNTf<sub>2</sub>, as initiators (10 mol %).<sup>25</sup> To our delight, we obtained in both cases the corresponding allylsilylated enamide **3a**, in 50% and 35% yield, respectively, with complete regio- and stereoselectivity (Table 1, entries 1 and 2).

**Table 1. Optimization of the Reaction of Ynamide **1a** and Allylsilane **2a**<sup>a</sup>**




entry	initiator (mol %)	<b>2a</b> (equiv)	solvent	yield <sup>b</sup> (%)
1	TMSNTf <sub>2</sub> (10)	2	DCE	50
2	HNTf <sub>2</sub> (10)	2	DCE	35
3	TMSOTf (10)	2	DCE	
4	TMSNTf <sub>2</sub> (10)	4	DCE	54
5	HNTf <sub>2</sub> (10)	4	DCE	65
6	TMSNTf <sub>2</sub> (5)	4	DCE	56
7	HNTf <sub>2</sub> (5)	4	DCE	66, 71 <sup>c</sup>
8	HNTf <sub>2</sub> (5)	4	DCM	64
9	HNTf <sub>2</sub> (5)	4	Et <sub>2</sub> O	16
10	HNTf <sub>2</sub> (5)	4	toluene	28
11	HNTf <sub>2</sub> (5)	4	THF	
12	HNTf <sub>2</sub> (5)	4	CH <sub>3</sub> CN	

<sup>a</sup>**1a** (0.1 mmol, 1 equiv), **2a** (equiv), initiator (mol %), solvent (0.4 M). <sup>b</sup>Isolated yield after flash chromatography purification on silica gel. <sup>c</sup>2 mmol scale.

We employed also trimethylsilyl trifluoromethanesulfonate, TMSOTf, as initiator in the same reaction conditions, but in this case, we observed only decomposition of the reagents (Table 1, entry 3). Then, the yields were improved by increasing the ratio of allylsilane **2a** to 4-fold excess (Table 1, entries 4,5); additionally, we observed that lowering the initiator loading to 5 mol % did not seem to affect the efficiency of the reaction (Table 1, entries 6 and 7). Furthermore, comparable yields were obtained with other halogenated solvents such as dichloromethane (Table 1, entry 8); however, the yields were significantly lower with diethyl ether or toluene (Table 1 entries 9 and 10), or simply the reaction did not afford any product when using THF or acetonitrile (Table 1, entries 11 and 12). Finally, the reaction was scaled-up to 2 mmol employing 5 mol % HNTf<sub>2</sub> with an improved yield of 71% (Table 1, entry 7).

With the optimized conditions in hand, we examined the scope of the reaction. We employed either TMSNTf<sub>2</sub> (method A) or HNTf<sub>2</sub> (method B) as initiators with different  $\beta$ -aryl-*N*-methyl-*N*-tosylamides **1** and allylsilane **2a** (Scheme 1).

**Scheme 1. Reaction of Ynamides **1** and Allyltrimethylsilane **2a****




<sup>a</sup>10 mol % initiator, DCE (0.2 M).

Thus,  $\alpha$ -allyl- $\beta$ -silyl-(*Z*)-enamides **3** were obtained with complete regio- and stereoselectivity and good yields (56–77%). In this way, it was proved that the reaction works well with different aryl-substituted ynamides bearing either electron-withdrawing (R = F, Br, CF<sub>3</sub>, products **3b–d,g,h**) or electron-donating (R = Me, MeO, products **3e,f**) substituents at different positions of the aryl moiety.

Then, we explored the reactivity of different 2-substituted allylsilanes **2** with a variety of  $\beta$ -aryl-substituted ynamides **1** (Scheme 2). The results were similar or even better than those previously obtained for the parent allyltrimethylsilane **2a** (R<sup>2</sup> = H). Thus, trimethyl(2-methylallyl)silane **2b** gave good yields with diverse ynamides (Scheme 2, R<sup>2</sup> = Me, products **3i–n**); furthermore, other 2-substituted allylsilanes like trimethyl-(2-phenylallyl)silane **2c** gave also excellent results (Scheme 2, R<sup>2</sup> = Ph, products **3o–u**). Interestingly, product **3o** was obtained almost quantitatively (99%) when the reaction was performed in a 2 mmol scale (Scheme 2). In addition, the structures of products **3j** and **3q** were unambiguously confirmed by X-ray resolution.<sup>26</sup> Continuing our study on 2-arylsubstituted allylsilanes, [2-(4-chlorophenyl)allyl]trimethylsilane **2d** (R<sup>2</sup> = 4-ClC<sub>6</sub>H<sub>4</sub>) and [2-(4-*t*-butylphenyl)allyl]trimethylsilane **2e** (R<sup>2</sup> = <sup>t</sup>BuC<sub>6</sub>H<sub>4</sub>) were also employed to obtain the corresponding silylenamides again with good yields (Scheme 2 products **3v,w** and **3x**, respectively). Regarding other substitution in the ynamide, we also checked a  $\beta$ -alkyl tosylamide (R<sup>1</sup> = *n*-butyl) with allylsilanes **2a,b** (R<sup>2</sup> = H, Me) in similar reaction conditions, but in this case only complex mixtures were obtained.<sup>27</sup> However, when an alkenyl  $\beta$ -substituted tosylamide (Scheme 2, R<sup>1</sup> = cinnamyl) was reacted in the same reaction conditions with allylsilane **2c** (R<sup>2</sup> = Ph), the expected  $\beta$ -silyl-(*Z*)-enamide **3y** was obtained in 51% yield (Scheme 2).

To get some insight into the reaction mechanism, we performed computational studies at the PCM-M05-2X/6-31G\*\*/M05-2X/6-31G\* level.<sup>26</sup> Starting from tosylamide **1a** and TMSNTf<sub>2</sub>, the molecular geometry was fully optimized without any molecular symmetry constraint, leading to structure **A** (Figure 2), a coordination minimum that placed the Si–C<sub>ph</sub> distance at 4.065 Å, keeping the Si–N bond distance at 1.895 Å. Subsequently, the approach between Si and C<sub>ph</sub> (Si–C<sub>ph</sub> distance = 2.195 Å) induces an elongation

**Scheme 2. Reaction of Ynamides **1** with 2-Substituted Allyltrimethylsilanes **2****


<sup>a</sup>10 mol % initiator, DCE (0.2 M). <sup>b</sup>99% 2 mol scale, method B. <sup>c</sup>Ellipsoids at 50% of probability level.


between Si and N (Si–N distance = 2.538 Å), giving rise to transition state **TS1** (+10.3 kcal·mol<sup>-1</sup>), in which the SiMe<sub>3</sub> moiety is rather flat (C–Si–C–C dihedral angle = 165.4°). **TS1** evolves to **B** (+6.2 kcal·mol<sup>-1</sup>), with formation of the Si–C<sub>Ph</sub> bond (Si–C<sub>Ph</sub> distance = 2.058 Å) and cleavage of the Si–N bond (Si–N distance = 2.906 Å). As also shown in **Figure 2**, the *anti*-approximation of allyltrimethylsilane **2a** to **B** gave the coordination minimum **C<sub>anti</sub>** (+0.4 kcal·mol<sup>-1</sup>), with a distance

of 4.093 Å between H<sub>2</sub>C= and C<sub>N</sub>, which is reduced in the transition state **TS2<sub>anti</sub>** (+8.8 kcal·mol<sup>-1</sup>) to 2.048 Å. **TS2<sub>anti</sub>** led to the minimum **D<sub>anti</sub>** (–1.2 kcal·mol<sup>-1</sup>), which in fact has a cyclopropyl structure (bond distances: H<sub>2</sub>C–C<sub>N</sub> = 1.592 Å, HC–C<sub>N</sub> = 1.565 Å, and H<sub>2</sub>C–HC = 1.464 Å). Finally, the attack of the Tf<sub>2</sub>N<sup>-</sup> anion on the silicon atom acts as the driving force of the process, leading directly to the coordination minimum **E<sub>anti</sub>** (–41.4 kcal·mol<sup>-1</sup>), formed by the allylsilylated enamide **3a** and TMSNTf<sub>2</sub>, without any intermediate being located. Likewise, *syn*-addition from the coordination minimum **C<sub>syn</sub>** (+2.1 kcal·mol<sup>-1</sup>) leads to the minimum **D<sub>syn</sub>** (–2.9 kcal·mol<sup>-1</sup>) through the transition state **TS2<sub>syn</sub>** (+13.6 kcal·mol<sup>-1</sup>), which shows a distance of 2.300 Å between H<sub>2</sub>C= and C<sub>N</sub>. The difference in the energy barriers to reach **TS2<sub>anti</sub>** or **TS2<sub>syn</sub>** (4.8 kcal·mol<sup>-1</sup>) could be explained by the β-silicon effect,<sup>19</sup> which places the bond angles in **B** at 107° (Si–C<sub>Ph</sub>–C<sub>N</sub>) and 131° (Ph–C<sub>Ph</sub>–C<sub>N</sub>), which avoid *syn*-approximation and allow us to explain the experimentally found stereoselectivity.<sup>26</sup>

According to our calculations, the overall process would start from the reaction of **1a** with TMSNTf<sub>2</sub> and formation of a β-silyl ketenimium intermediate **B** and the Tf<sub>2</sub>N<sup>-</sup> anion. Intermediate **B** would receive nucleophilic *anti*-attack of the C–C double bond of the allylic silane **2a** to give the intermediate **D<sub>anti</sub>**. The subsequent attack of the Tf<sub>2</sub>N<sup>-</sup> anion on the silicon atom leads to β-silylenamide **3a** and TMSNTf<sub>2</sub>, which closes the catalytic cycle (**Figure 2**).


Finally, to illustrate the synthetic possibilities of enamides **3**, we carried out several transformations (**Scheme 3**). Thus, the reaction of β-silylenamide **3o** with a fluoride source, such as tetrabutylammonium fluoride, led to the desilylated enamide **4** (86%) or, alternatively, the coupling product **5** (72%) if the reaction was performed in the presence of 4-bromobenzaldehyde (**Scheme 3**). The allyl group can also intervene in other transformations; thus, the presence of a catalytic amount of a Brønsted acid (HNTf<sub>2</sub>, 1 mol %) gave the 1,2-dihydronaphthalene derivative **6** (65%) because of an intramolecular aromatic electrophilic substitution of the carbocation intermediate formed by previous protonation of the 2-phenylallyl substituent (**Scheme 3**).

In summary, we have described a regio- and stereoselective carbosilylation of tosylynamides **1** catalyzed by silylium ion. The reaction uses different allylsilanes **2** as the source of the



**Figure 2.** Calculated relative energy profile for the formation of β-silylenamides, in kcal·mol<sup>-1</sup> (for the sake of comparison, the values shown for **A**, **TS1**, and **B** also include the energy value of **2a**) and overall reaction of the proposed catalytic cycle. H atoms have been omitted for clarity.

## Scheme 3. Further Transformations of silylenamide 3o



carbon nucleophile and the silicon electrophile. The silylium ion activates the triple C–C bond of the ynamide to produce an electrophilic  $\beta$ -silylketenimonium intermediate **B**, the subsequent nucleophilic attack by the allylsilane **2** produces the regeneration of the silylium ion to close the catalytic cycle. Theoretical calculations support this mechanistic picture. This versatile reaction leads to (*Z*)- $\beta$ -silylenamides **3**, interesting building blocks as demonstrated by the possibility of further transformations. Finally, this reaction represents a novel example of catalytic activation of electron rich alkynes by silylium ion.

### ■ ASSOCIATED CONTENT

#### Data Availability Statement

The data underlying this study are available in the published article and its online [Supporting Information](#).

#### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.3c00221>.

Experimental details, materials and methods, characterization data, NMR spectra for all compounds, X-ray diffraction experiments and computational studies data ([PDF](#))

#### Accession Codes

CCDC 2193409–2193410 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

### ■ AUTHOR INFORMATION

#### Corresponding Author

Alfredo Ballesteros – *Departamento de Química Orgánica e Inorgánica, Instituto Universitario de Química Organometálica “Enrique Moles”, Universidad de Oviedo, 33006 Oviedo, Spain*; [orcid.org/0000-0003-2093-4444](https://orcid.org/0000-0003-2093-4444); Email: [abg@uniovi.es](mailto:abg@uniovi.es)

#### Authors

Paz Yepes – *Departamento de Química Orgánica e Inorgánica, Instituto Universitario de Química Organometálica “Enrique Moles”, Universidad de Oviedo, 33006 Oviedo, Spain*  
 Ángel L. Suárez-Sobrino – *Departamento de Química Orgánica e Inorgánica, Instituto Universitario de Química*

*Organometálica “Enrique Moles”, Universidad de Oviedo, 33006 Oviedo, Spain*

Miguel A. Rodríguez – *Departamento de Química, Centro de Investigación en Síntesis Orgánica, Universidad de la Rioja, 26006 Logroño, Spain*

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.orglett.3c00221>

### Notes

The authors declare no competing financial interest.

### ■ ACKNOWLEDGMENTS

This research was supported by grant from AEI (PID2019-107469RB-I00/AEI/10.13039/501100011033). The authors also thank Prof. J. M. González (Universidad de Oviedo) for helpful discussions and suggestions. Computing resources used in this work were provided by Universidad de La Rioja (Beronia cluster). The authors also acknowledge the technical support provided by *Servicios Científico-Técnicos* de la Universidad de Oviedo.

### ■ DEDICATION

Dedicated to the memory of Professor Gregorio Asensio.

### ■ REFERENCES

- (1) (a) Shaykhtudinova, P.; Keess, S.; Oestreich, M. in *Organosilicon Chemistry-Novel Approaches and Reactions*; Hiyama, T., Oestreich, M., Eds.; Wiley-VCH, Weinheim, 2019; pp 131–170. (b) Klare, H. F. T.; Albers, L.; Süsse, L.; Keess, S.; Müller, T.; Oestreich, M. Silylium Ions: From Elusive Reactive Intermediates to Potent Catalyst. *Chem. Rev.* **2021**, *121*, 5889–5985.
- (2) Lambert, J. B.; Zhao, Y.; Wu, H.  $\beta$ -Silyl and  $\beta$ -Germlyl Carbocations Stable at Room Temperature. *J. Org. Chem.* **1999**, *64*, 2729–2736.
- (3) Intramolecular hydrosilylation of alkenes: Steinberger, H.-U.; Bauch, C.; Müller, T.; Auner, N. A. A metal-free catalytic intramolecular hydrosilylation. *Can. J. Chem.* **2003**, *81*, 1223–1227.
- (4) Intermolecular hydrosilylation of alkenes intercepted by a Friedel–Crafts reaction: Arii, H.; Yano, Y.; Nakabayashi, K.; Yamaguchi, S.; Yamamura, M.; Mochida, K.; Kawashima, T. Regioselective and Stereospecific Dehydrogenerative Annulation Utilizing Silylium Ion-Activated Alkenes. *J. Org. Chem.* **2016**, *81*, 6314–6319.
- (5) Reactions of vinylcyclopropanes and hydrosilanes promoted by silylium ion: He, T.; Wang, G.; Long, P.-W.; Kemper, S.; Irran, E.; Klare, H. F. T.; Oestreich, M. Intramolecular Friedel–Crafts alkylation with a silylium-ion-activated cyclopropyl group: formation of tricyclic ring systems from benzyl-substituted vinylcyclopropanes and hydrosilanes. *Chem. Sci.* **2021**, *12*, 569–575 and references therein.
- (6) Rej, S.; Klare, H. F. T.; Oestreich, M. Silylium-Ion-Promoted Hydrosilylation of Aryl-Substituted Allenes: Interception by Cyclization of the Allyl-Cation Intermediate. *Org. Lett.* **2022**, *24*, 1346–1350.
- (7) He, T.; Qu, Z.-W.; Klare, H. F. T.; Grimme, S.; Oestreich, M. Intermolecular Carbosilylation of  $\alpha$ -Olefins with C(sp<sup>3</sup>)-C(sp) Bond Formation Involving Silylium-Ion Regeneration. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202203347.
- (8) (a) Müller, T.; Meyer, R.; Lennartz, D.; Siehl, H.-U. Unusually Stable Vinyl Cation. *Angew. Chem., Int. Ed.* **2000**, *39*, 3074–3077. (b) Martens, A.; Kreuzer, M.; Ripp, A.; Schneider, M.; Himmel, D.; Scherer, H.; Krossing, I. Investigations on non-classical silylium ions leading to a cyclobutenyl cation. *Chem. Sci.* **2019**, *10*, 2821–2829.
- (9) Arii, H.; Kurihara, T.; Mochida, K.; Kawashima, T. Silylium ion-promoted dehydrogenerative cyclization synthesis of silicon-containing compounds derived from alkynes. *Chem. Commun.* **2014**, *50*, 6649–6652.

(10) (a) Chen, Y.-B.; Qian, P.-C.; Ye, L.-W. Brønsted acid-mediated reactions of ynamides. *Chem. Soc. Rev.* **2020**, *49*, 8897–8909. (b) Hong, F.-L.; Ye, L.-W. Transition Metal Catalyzed Tandem Reactions of Ynamides for Divergent *N*-Heterocycle Synthesis. *Acc. Chem. Res.* **2020**, *53*, 2003–2019. (c) Campeau, D.; Rayo, D. F. L.; Mansour, A.; Muratov, K.; Gagosz, F. Gold-Catalyzed Reactions of Specially Activated Alkynes, Allenes, and Alkenes. *Chem. Rev.* **2021**, *121*, 8756–8867.

(11) (a) Evano, G.; Coste, A.; Jouvin, K. Ynamides: Versatile Tools in Organic Synthesis. *Angew. Chem., Int. Ed.* **2010**, *49*, 2840–2859. (b) Dodd, R. H.; Cariou, K. Ketanimines Generated from Ynamides: Versatile Building Blocks for Nitrogen-Containing Scaffolds. *Chem.—Eur. J.* **2018**, *24*, 2297–2304.

(12) Recent examples: (a) Takimoto, M.; Gholap, S. S.; Hou, Z. Alkylative Carboxylation of Ynamides and Allenamides with Functionalized Alkylzinc Halides and Carbon Dioxide by a Copper Catalyst. *Chem.—Eur. J.* **2019**, *25*, 8363–8370. (b) Dutta, S.; Yang, S.; Vanjari, R.; Mallick, R. K.; Gandon, V.; Sahoo, A. K. Keteniminium-Driven Umpolung Difunctionalization of Ynamides. *Angew. Chem., Int. Ed.* **2020**, *59*, 10785–10790. (c) You, C.; Sakai, M.; Daniliuc, C. G.; Bergander, K.; Yamaguchi, S.; Studer, A. Regio- and Stereoselective 1,2-Carbaboration of Ynamides with Aryldichloroboranes. *Angew. Chem., Int. Ed.* **2021**, *60*, 21697–21701.

(13) Szudkowska-Frątczak, J.; Hreczycho, G.; Pawluć, P. Silylative coupling of olefins with vinylsilanes in the synthesis of functionalized alkenes. *Org. Chem. Front.* **2015**, *2*, 730–738.

(14) Vercrussse, S.; Jouvin, K.; Riant, O.; Evano, G. Copper-Catalyzed Silylcupration of Activated Alkynes. *Synthesis* **2016**, *48*, 3373–3381.

(15) Timbart, L.; Cintrat, J.-C. Synthesis and Reactivity of an  $\alpha$ -Stannyl  $\beta$ -Silyl Enamide. *Chem.—Eur. J.* **2002**, *8*, 1637–1640 and references therein.

(16) Saito, N.; Saito, K.; Sato, H.; Sato, Y. Regio- and Stereoselective Synthesis of Tri- and Tetrasubstituted Enamides via Palladium-Catalyzed Silaboration of Ynamides. *Adv. Synth. Catal.* **2013**, *355*, 853–857.

(17) (a) Romain, E.; Fopp, C.; Chemla, F.; Ferreira, F.; Jackowski, O.; Oestreich, M.; Pérez-Luna, A. *Trans*-Selective Radical Silylzincation of Ynamides. *Angew. Chem., Int. Ed.* **2014**, *53*, 11333–11337. (b) A stereodivergent silylzincation of ynamides giving rise selectively to *Z* or *E*  $\beta$ -silylenamides was later described by the same group: Fopp, C.; Romain, E.; Isaac, K.; Chemla, F.; Ferreira, F.; Jackowski, O.; Oestreich, M.; Pérez-Luna, A. Stereodivergent Silylzincation of  $\alpha$ -Heteroatom-Substituted Alkynes. *Org. Lett.* **2016**, *18*, 2054–2057.

(18) Zheng, N.; Song, W.; Zhang, T.; Li, M.; Zheng, Y.; Chen, L. Rhodium-Catalyzed Highly Regioselective and Stereoselective Intermolecular Hydrosilylation of Internal Ynamides under Mild Conditions. *J. Org. Chem.* **2018**, *83*, 6210–6216.

(19) Kim, Y.; Dateer, R. B.; Chang, S. Borane-Catalyzed Selective Hydrosilylation of Internal Ynamides Leading to  $\beta$ -Silyl (*Z*)-Enamides. *Org. Lett.* **2017**, *19*, 190–193.

(20) Hansjacob, P.; Leroux, F. R.; Gandon, V.; Donnard, M. Palladium-Catalyzed Silylcyanation of Ynamides: Regio- and Stereoselective Access to Tetrasubstituted 3-Silyl-2-Aminoacrylonitriles. *Angew. Chem. Int. Ed.* **2022**, *61*, No. e202200204.

(21) Allylsilylation of alkenes: Motokura, K.; Matsunaga, S.; Miyaji, A.; Sakamoto, Y.; Baba, T. Heterogeneous Allylsilylation of Aromatic and Aliphatic Alkenes Catalyzed by Proton-Exchanged Montmorillonite. *Org. Lett.* **2010**, *12*, 1508–1511.

(22) Allylsilylation of alkynes: Imamura, K.-I.; Yoshikawa, E.; Gevorgyan, V.; Yamamoto, Y. First Exclusive *Endo-dig* Carbocyclization: HfCl<sub>4</sub>-Catalyzed Intramolecular Allylsilylation of Alkynes. *J. Am. Chem. Soc.* **1998**, *120*, 5339–5340 and references therein.

(23) For a review: Motokura, K.; Baba, T. An atom-efficient synthetic method: carbosilylations of alkenes, alkynes, and cyclic acetals using Lewis and Brønsted acid catalysis. *Green Chem.* **2012**, *14*, 565–579.

(24) Rubial, B.; Ballesteros, A.; González, J. M. Silylium-Catalyzed Carbon–Carbon Coupling of Alkynylsilanes with (2-Bromo-1-

methoxyethyl)arenes: Alternative Approaches. *Eur. J. Org. Chem.* **2018**, *2018*, 6194–6198 and references therein.

(25) Rubial, B.; Ballesteros, A.; González, J. M. Silylium-Catalyzed Alkynylation and Etherification Reactions of Benzylic Acetates. *Eur. J. Org. Chem.* **2022**, No. e202200051, and references therein.

(26) See the [Supporting Information](#) for additional details.

(27) The presence of acidic hydrogens in the alkyl moiety opens the possibility of other transformations of silylketeniminium cation intermediate **B** in the same reaction conditions.

## Recommended by ACS

### Modular Synthesis of (Borylmethyl)silanes through Orthogonal Functionalization of a Carbon Atom

Rajdip Chowdhury, Abraham Mendoza, *et al.*

MARCH 10, 2023

ORGANIC LETTERS

[READ](#)

### Catalytically Generated Meerwein's Salt-Type Oxonium Ions for Friedel–Crafts C(sp<sup>2</sup>)–H Methylation with Methanol

Tao He, Martin Oestreich, *et al.*

FEBRUARY 01, 2023

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

[READ](#)

### $\beta$ -Boron Effect Enables Regioselective and Stereospecific Electrophilic Addition to Alkenes

Yin Li, Andrei K. Yudin, *et al.*

MARCH 22, 2023

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

[READ](#)

### Gold-Catalyzed Regio- and Stereoselective $\alpha$ -Acyloxy- $\beta$ -Alkynylation of Ynol Ethers

Yaowen Liu, A. Stephen K. Hashmi, *et al.*

FEBRUARY 17, 2023

THE JOURNAL OF ORGANIC CHEMISTRY

[READ](#)

[Get More Suggestions >](#)